

## Lab efficacy

- Altreuther, G., Schimmel, A., Schroeder, I., Bach, T., Charles, S., Kok, D. J., Krieger, K. J. (2009). Efficacy of emodepside plus praziquantel tablets (Profender tablets for dogs) against mature and immature infections with *Toxocara canis* and *Toxascaris leonina* in dogs. *Parasitol Res*, 105 Suppl 1, S1-8. doi: 10.1007/s00436-009-1489-7
- Baker, C., Tielemans, E., Prullage, J. B., Chester, S. T., Knaus, M., Rehbein, S., Rosentel, J. K. (2014). Efficacy of a novel topical combination of fipronil, (S)-methoprene, eprinomectin and praziquantel against adult and immature stages of the cat flea (*Ctenocephalides felis*) on cats. *Veterinary Parasitology*, 202(1–2), 54-58. doi: <http://dx.doi.org/10.1016/j.vetpar.2014.02.040>
- Beugnet, F., Delpont, P., Luus, H., Crafford, D., & Fourie, J. (2013). Preventive efficacy of Frontline((R)) Combo and Certifect((R)) against *Dipylidium caninum* infestation of cats and dogs using a natural flea (*Ctenocephalides felis*) infestation model. *Parasite*, 20, 7. doi: 10.1051/parasite/2013006
- Beugnet, F., deVos, C., Liebenberg, J., Halos, L., & Fourie, J. (2014). Afoxolaner against fleas: immediate efficacy and resultant mortality after short exposure on dogs. *Parasite*, 21, 42. doi: 10.1051/parasite/2014045
- Beugnet, F., Fourie, J., & Chalvet-Monfray, K. (2012). Comparative efficacy on dogs of a single topical treatment with fipronil/(S)-methoprene or weekly physiological hygiene shampoos against *Ctenocephalides felis* in a simulated flea-infested environment. *Parasite: journal de la Société Française de Parasitologie*, 19(2), 153.
- Beugnet, F., Halos, L., Larsen, D., Labuschagné, M., Erasmus, H., & Fourie, J. (2014). The ability of an oral formulation of afoxolaner to block the transmission of *Babesia canis* by *Dermacentor reticulatus* ticks to dogs. *Parasites & Vectors*, 7(1), 1-7. doi: 10.1186/1756-3305-7-283
- Beugnet, F., Liebenberg, J., & Halos, L. (2015). Comparative efficacy of two oral treatments for dogs containing either afoxolaner or fluralaner against *Rhipicephalus sanguineus sensu lato* and *Dermacentor reticulatus*. *Vet Parasitol*. doi: 10.1016/j.vetpar.2015.02.002
- Beugnet, F., Liebenberg, J., & Halos, L. (2015). Comparative speed of efficacy against *Ctenocephalides felis* of two oral treatments for dogs containing either afoxolaner or fluralaner. *Vet Parasitol*, 207(3-4), 297-301. doi: 10.1016/j.vetpar.2014.12.007
- Bienhoff, S. E., Kok, D. J., Roycroft, L. M., & Roberts, E. S. (2013). Efficacy of a single oral administration of milbemycin oxime against natural infections of *Ancylostoma braziliense* in dogs. *Vet Parasitol*. doi: 10.1016/j.vetpar.2013.01.004
- Bonneau, S., Maynard, L., Tomczuk, K., Kok, D., & Eun, H. M. (2009). Anthelmintic efficacies of a tablet formula of ivermectin-praziquantel on horses experimentally infected with three *Strongylus* species. *Parasitol Res*, 105(3), 817-823. doi: 10.1007/s00436-009-1460-7
- Cadiergues, M. C., Bonneau, S. G., & Fourier, J. J. (2011). Assay of two 10% (w/v) fipronil spot-on formulations against feline infestations with *Ctenocephalides felis*. *J Feline Med Surg*, 13(4), 304-308. doi: 10.1016/j.jfms.2011.01.004
- Charles, S. D., Altreuther, G., Reinemeyer, C. R., Buch, J., Settje, T., Cruthers, L., Kok, D. J., Bowman, D. D., Kazacos, K. R., Jenkins, D. J. & Schein, E. (2005). Evaluation of the efficacy of emodepside+praziquantel topical solution against cestode (*Dipylidium caninum*, *Taenia taeniaeformis*, and *Echinococcus multilocularis*) infections in cats. *Parasitol Res*, 97 Suppl 1, S33-40. doi: 10.1007/s00436-005-1442-3
- Dumont, P., Blair, J., Fourie, J. J., Chester, T. S., & Larsen, D. L. (2014). Evaluation of the efficacy of afoxolaner against two European dog tick species: *Dermacentor reticulatus* and *Ixodes ricinus*. *Veterinary Parasitology*, 201(3–4), 216-219. doi: <http://dx.doi.org/10.1016/j.vetpar.2014.02.017>
- Fankhauser, B., Dumont, P., Halos, L., Hunter, J. S., Kunkle, B., Everett, W. R., Chester, T. S., Fourie, J. J. & Soll, M. (2015). Efficacy of a new combination of fipronil and permethrin against *Ctenocephalides felis* flea infestation in dogs. *Parasit Vectors*, 8(1), 62. doi: 10.1186/s13071-015-0687-7

- Fourie, J. J., Beugnet, F., Ollagnier, C., & Pollmeier, M. G. (2011). Study of the sustained speed of kill of the combination of fipronil/amitraz/(S)-methoprene and the combination of imidacloprid/permethrin against *Dermacentor reticulatus*, the European dog tick. *Parasite*, 18(4), 319-323.
- Fourie, J. J., Crafford, D., Horak, I. G., & Stanneck, D. (2012). Prophylactic treatment of flea-infested cats with an imidacloprid/flumethrin collar to forestall infection with *Dipylidium caninum*. *Parasit Vectors*, 5, 151.
- Fourie, J. J., Crafford, D., Horak, I. G., & Stanneck, D. (2013). Prophylactic treatment of flea-infested dogs with an imidacloprid / flumethrin collar (Seresto(R), Bayer) to preempt infection with *Dipylidium caninum*. *Parasitol Res*, 112 Suppl 1, 33-46. doi: 10.1007/s00436-013-3279-5
- Fourie, J. J., Fourie, L. J., & Horak, I. G. (2005). Efficacy of orally administered powdered aloe juice (*Aloe ferox*) against ticks on cattle and ticks and fleas on dogs. *J S Afr Vet Assoc*, 76(4), 193-196.
- Fourie, J. J., Fourie, L. J., Horak, I. G., & Snyman, M. G. (2010). The efficacy of a topically applied combination of cyphenothrin and pyriproxyfen against the southern African yellow dog tick, *Haemaphysalis elliptica*, and the cat flea, *Ctenocephalides felis*, on dogs. *J S Afr Vet Assoc*, 81(1), 33-36.
- Fourie, J. J., Horak, I., Crafford, D., Erasmus, H. L., & Botha, O. J. (2015). The efficacy of a generic doxycycline tablet in the treatment of canine monocytic ehrlichiosis (Vol. 86).
- Fourie, J. J., Luus, H. G., Stanneck, D., & Jongejan, F. (2013). The efficacy of Advantix(R) to prevent transmission of *Ehrlichia canis* to dogs by *Rhipicephalus sanguineus* ticks. *Parasite*, 20, 36. doi: 10.1051/parasite/2013037
- Fourie, J. J., Ollagnier, C., Beugnet, F., Luus, H. G., & Jongejan, F. (2013). Prevention of transmission of *Ehrlichia canis* by *Rhipicephalus sanguineus* ticks to dogs treated with a combination of fipronil, amitraz and (S)-methoprene (CERTIFECT((R))). *Vet Parasitol*, 193(1-3), 223-228. doi: 10.1016/j.vetpar.2012.12.009
- Fourie, J. J., Stanneck, D., & Jongejan, F. (2013). Prevention of transmission of *Babesia canis* by *Dermacentor reticulatus* ticks to dogs treated with an imidacloprid/flumethrin collar. *Veterinary Parasitology*, 192(1-3), 273-278. doi: <http://dx.doi.org/10.1016/j.vetpar.2012.10.017>
- Fourie, J. J., Stanneck, D., Luus, H. G., Beugnet, F., Wijnveld, M., & Jongejan, F. (2013). Transmission of *Ehrlichia canis* by *Rhipicephalus sanguineus* ticks feeding on dogs and on artificial membranes. *Veterinary Parasitology*, 197(3-4), 595-603. doi: <http://dx.doi.org/10.1016/j.vetpar.2013.07.026>
- Fourie, L. J., Kok, D. J., Allan, M. J., & Oberem, P. T. (1995). The efficacy of Diflubenzuron against the body louse (*Damalinea limbata*) of Angora goats. *Vet Parasitol*, 59(3-4), 257-262.
- Fourie, L. J., Kok, D. J., & Heine, J. (2003). Evaluation of the efficacy of an imidacloprid 10%/moxidectin 1% spot-on against *Otodectes cynotis* on cats. *Parasitol Res*, 90 Suppl 3, S112-113. doi: 10.1007/s00436-003-0906-6
- Fourie, L. J., Kok, D. J., & Peter, R. J. (2000). Control of immature stages of the flea *Ctenocephalides felis* (Bouché) in carpets exposed to cats treated with imidacloprid. *J S Afr Vet Assoc*, 71(4), 219-221.
- Fourie, L. J., Stanneck, D., & Horak, I. G. (2003). The efficacy of collars impregnated with flumethrin and propoxur against experimental infestations of adult *Rhipicephalus sanguineus* on dogs. *J S Afr Vet Assoc*, 74(4), 123-126.
- Fourie, L. J., Stanneck, P. D., & Horak, I. G. (2006). The Efficacy of a Topically Applied Combination of Imidacloprid and Permethrin Against *Stomoxys calcitrans* on Dogs. *Intern J Appl Res Vet Med*, 4(1), 29-33.
- Horak, I. G., Fourie, J. J., & Stanneck, D. (2012). Efficacy of slow-release collar formulations of imidacloprid/flumethrin and deltamethrin and of spot-on formulations of fipronil/(s)-methoprene, dinotefuran/pyriproxyfen/permethrin and (s)-methoprene/amitraz/fipronil against *Rhipicephalus sanguineus* and *Ctenocephalides felis felis* on dogs. *Parasit Vectors*, 5, 79. doi: 10.1186/1756-3305-5-79
- Hunter Iii, J. S., Baggott, D., Everett, W. R., Fourie, J. J., Cramer, L. G., Yoon, S. S., Collidor, N., Mallouk, Y., Lee, L., Blair, J. & Prullage, J. B. (2011). Efficacy of a novel topical combination of

fipronil, amitraz and (S)-methoprene for treatment and control of induced infestations of brown dog ticks (*Rhipicephalus sanguineus*) on dogs. *Veterinary Parasitology*, 179(4), 318-323. doi: <http://dx.doi.org/10.1016/j.vetpar.2011.03.042>

Jongejan, F., de Vos, C., Fourie, J. J., & Beugnet, F. (2015). A novel combination of fipronil and permethrin (Frontline Tri-Act(R)/Frontect(R)) reduces risk of transmission of *Babesia canis* by *Dermacentor reticulatus* and of *Ehrlichia canis* by *Rhipicephalus sanguineus* ticks to dogs. *Parasit Vectors*, 8, 602. doi: 10.1186/s13071-015-1207-5

Jongejan, F., Fourie, J. J., Chester, S. T., Manavella, C., Mallouk, Y., Pollmeier, M. G., & Baggott, D. (2011). The prevention of transmission of *Babesia canis canis* by *Dermacentor reticulatus* ticks to dogs using a novel combination of fipronil, amitraz and (S)-methoprene. *Vet Parasitol*, 179(4), 343-350. doi: 10.1016/j.vetpar.2011.03.047

King, J. N., Delpont, P. C., Luus, H. G., Erasmus, H. L., Barnes, P. M., & Speranza, C. (2014). Efficacy and acceptability of the new oral phosphate binder Lenziaren® in healthy cats fed a renal diet. *Journal of Veterinary Pharmacology and Therapeutics*, n/a-n/a. doi: 10.1111/jvp.12173

King, J. N., Erasmus, H. L., Delpont, P. C., Bester, I. C., & Seewald, W. (2014). Efficacy, acceptability and tolerability of the new oral phosphate binder Lenziaren® in healthy cats fed a standard diet. *BMC Veterinary Research*, 10(1), 258.

Kok, D. J., Schenker, R., Archer, N. J., Horak, I. G., & Swart, P. (2011). The efficacy of milbemycin oxime against pre-adult *Spirocerca lupi* in experimentally infected dogs. *Veterinary Parasitology*, 177(1-2), 111-118. doi: <http://dx.doi.org/10.1016/j.vetpar.2010.11.031>

Kok, D. J., Williams, E. J., Schenker, R., Archer, N. J., & Horak, I. G. (2010). The use of milbemycin oxime in a prophylactic anthelmintic programme to protect puppies, raised in an endemic area, against infection with *Spirocerca lupi*. *Veterinary Parasitology*, 174(3-4), 277-284. doi: <http://dx.doi.org/10.1016/j.vetpar.2010.08.029>

Kuzner, J., Turk, S., Fourie, J. J., Grace, S., Marchiondo, A. A., & Rugg, D. (2013). Efficacy of a novel fipronil spot-on for the treatment and control of induced infestations of adult cat fleas (*Ctenocephalides felis*) and castor bean ticks (*Ixodes ricinus*) on cats. *Parasitol Res*, 112(1), 365-372. doi: 10.1007/s00436-012-3144-y

Kuzner, J., Turk, S., Grace, S., Soni-Gupta, J., Fourie, J. J., Marchiondo, A. A., & Rugg, D. (2013). Confirmation of the efficacy of a novel fipronil spot-on for the treatment and control of fleas, ticks and chewing lice on dogs. *Vet Parasitol*, 193(1-3), 245-251. doi: 10.1016/j.vetpar.2012.11.006

Navarro, C., Reymond, N., Fourie, J., Hellmann, K., & Bonneau, S. (2015). Prevention of *Babesia canis* in dogs: efficacy of a fixed combination of permethrin and fipronil (Effitix(R)) using an experimental transmission blocking model with infected *Dermacentor reticulatus* ticks. *Parasit Vectors*, 8(1), 32. doi: 10.1186/s13071-015-0645-4

Schimmel, A., Altreuther, G., Schroeder, I., Charles, S., Cruthers, L., Ketzis, J., Kraemer, F., McCall, J. W., & Krieger, K. J. (2009). Efficacy of emodepside plus praziquantel tablets (Profender tablets for dogs) against mature and immature adult *Ancylostoma caninum* and *Uncinaria stenocephala* infections in dogs. *Parasitol Res*, 105 Suppl 1, S9-16. doi: 10.1007/s00436-009-1490-1

Schroeder, I., Altreuther, G., Schimmel, A., Deplazes, P., Kok, D. J., Schnyder, M., & Krieger, K. J. (2009). Efficacy of emodepside plus praziquantel tablets (Profender tablets for dogs) against mature and immature cestode infections in dogs. *Parasitol Res*, 105 Suppl 1, S31-38. doi: 10.1007/s00436-009-1493-y

Schmidt, B., Delpont, P., & SPIECKER-HAUSER, U. (2006). G07 Bay 78-1887, a novel lanthanum-based phosphate binder, decreases intestinal phosphorus absorption in cats. *Journal of Veterinary Pharmacology and Therapeutics*, 29(s1), 206-207.

Schmidt, B. H., Dribusch, U., Delpont, P. C., Gropp, J. M., & van der Staay, F. J. (2012). Tolerability and efficacy of the intestinal phosphate binder Lantharenol(R) in cats. *BMC Vet Res*, 8, 14. doi: 10.1186/1746-6148-8-14

Taenzler, J., Wengenmayer, C., Williams, H., Fourie, J., Zschiesche, E., Roepke, R. K. A., & Heckerroth, A. R. (2014). Onset of activity of fluralaner (BRAVECTO™) against *Ctenocephalides felis* on dogs. *Parasites & Vectors*, 7(1), 567. doi: 10.1186/s13071-014-0567-6

Varloud, M., & Fourie, J. (2015). One-month comparative efficacy of three topical ectoparasiticides against adult brown dog ticks (*Rhipicephalus sanguineus sensu lato*) on mixed-bred dogs in controlled environment. *Parasitology Research*, 1-9. doi: 10.1007/s00436-015-4356-8

Stanneck, D., & Fourie, J. J. (2013). Imidacloprid 10 % / flumethrin 4.5 % collars (Seresto(R), Bayer) successfully prevent long-term transmission of *Ehrlichia canis* by infected *Rhipicephalus sanguineus* ticks to dogs. *Parasitol Res*, 112 Suppl 1, 21-32. doi: 10.1007/s00436-013-3278-6